

COORDINATION COMPOUNDS - I

7. Coordination compounds

1. **Difference between coordination compound and double bond:**

Coordination compound	Double salt
A coordination compound contains a central metal atom or ion surrounded by number of oppositely charged ions or neutral molecules. These ions or molecules re bonded to the metal atom or ion by a coordinate bond.	When two salts in stoichiometric ratio are crystallised together from their saturated solution they are called double salts
Example: $K_4[Fe(CN)_6]$	Example: $FeSO_4 \cdot (NH_4)_2SO_4 \cdot 6H_2O$ (Mohr's salt)
They do not dissociate into simple ions when dissolved in water.	They dissociate into simple ions when dissolved in water.

- Coordination entity:** A coordination entity constitutes a central metal atom or ion bonded to a fixed number of ions or molecules. Example: in $K_4[Fe(CN)_6]$, $[Fe(CN)_6]^{4-}$ represents coordination entity.
- Central atom or ion:** In a coordination entity, the atom/ion to which a fixed number of ions/groups are bound in a definite geometrical arrangement around it, is called the central atom or ion. E.g.: in $K_4[Fe(CN)_6]$, Fe^{2+} is the central metal ion.
- Ligands:** A molecule, ion or group that is bonded to the metal atom or ion in a complex or coordination compound by a coordinate bond is called ligand. It may be neutral, positively or negatively charged. Example: H_2O , CN^- , NO^+ etc.
- Donor atom:** An atom of the ligand attached directly to the metal is called the donor atom. For example: in the complex $K_4[Fe(CN)_6]$, carbon is a donor atom.
- Coordination number:** The coordination number (CN) of a metal ion in a complex can be defined as the number of ligand donor atoms to

which the metal is directly bonded. For example: in the complex $K_4[Fe(CN)_6]$, the coordination number of Fe is 6.

7. **Coordination sphere:** The central atom/ion and the ligands attached to it are enclosed in square bracket and is collectively termed as the coordination sphere. For example: in the complex $K_4[Fe(CN)_6]$, $[Fe(CN)_6]^{4-}$ is the coordination sphere.
8. **Counter ions:** The ions present outside the coordination sphere are called counter ions. For example: in the complex $K_4[Fe(CN)_6]$, K^+ is the counter ion.
9. **Coordination polyhedron:** The spatial arrangement of the ligand atoms which are directly attached to the central atom/ ion defines a coordination polyhedron about the central atom. The most common coordination polyhedra are octahedral, square planar and tetrahedral. For example: $[PtCl_4]^{2-}$ is square planar, $Ni(CO)_4$ is tetrahedral while $[Cu(NH_3)_6]^{3+}$ is octahedral.
10. **Charge on the complex ion:** The charge on the complex ion is equal to the algebraic sum of the charges on all the ligands coordinated to the central metal ion.
11. **Denticity:** The number of ligating (linking) atoms present in ligand is called denticity.
12. **Unidentate ligands:** The ligands whose only one donor atom is bonded to metal atom are called unidentate ligands. Example: H_2O , NH_3 , CO , CN^-
13. **Didentate ligands:** The ligands which contain two donor atoms or ions through which they are bonded to the metal ion. For example: ethylene diamine ($H_2NCH_2CH_2NH_2$) has two nitrogen atoms, oxalate ion $\left(\begin{array}{c} COO^- \\ | \\ COO^- \end{array} \right)$ has two oxygen atoms which can bind with the metal atom.
14. **Polydentate ligand:** When several donor atoms are present in a single ligand, the ligand is called polydentate ligand. For example: in $N(CH_2CH_2NH_2)_3$, the ligand is said to be polydentate. Ethylenediaminetetraacetate ion ($EDTA^{4-}$) is an important hexadentate

ligand. It can bind through two nitrogen and four oxygen atoms to a central metal ion.

15. **Chelate:** An inorganic metal complex in which there is a close ring of atoms caused by attachment of a ligand to a metal atom at two points. An example is the complex ion formed between ethylene diamine and cupric ion, $[\text{Cu}(\text{NH}_2\text{CH}_2\text{NH}_2)_2]^{2+}$.
16. **Ambidentate ligand:** Ligands which can ligate (link) through two different atoms present in it are called ambidentate ligand. Example: NO_2^- , and SCN^- . NO_2^- can link through N as well as O while SCN^- can link through S as well as N atom.
17. **Werner's coordination theory:** Werner was able to explain the nature of bonding in complexes. The postulates of Werner's theory are:
 - a. Metal shows two different kinds of valencies: primary valence and secondary valence.

Primary valence	Secondary valence
This valence is normally ionisable.	This valence is non – ionisable.
It is equal to positive charge on central metal atom.	The secondary valency equals the number of ligand atoms coordinated to the metal. It is also called coordination number of the metal.
These valencies are satisfied by negatively charged ions.	It is commonly satisfied by neutral and negatively charged, sometimes by positively charged ligands.
Example: in CrCl_3 , the primary valency is three. It is equal to oxidation state of central metal ion.	

- b. The ions/ groups bound by secondary linkages to the metal have characteristic spatial arrangements corresponding to different coordination numbers.
- c. The most common geometrical shapes in coordination compounds are octahedral square planar and tetrahedral